

**Service Level Agreement
Basic Service: Consent
Version 1.2**

This document is provided to you free of charge by the

eHealth platform

**Willebroekkaai 38 – 1000 Brussel
38, Quai de Willebroeck – 1000 Bruxelles**

All are free to circulate this document with reference to the URL source.

Table of contents

Table of contents	2
1. Document management	4
1.1 Document history	4
1.2 Document references	4
1.3 Purpose of the document	4
1.4 Features	4
1.5 Validity of the agreement	5
1.6 Service and maintenance window	5
1.6.1 Service window	5
1.6.2 Support window	6
1.6.3 Maintenance windows & planned interventions	6
1.6.4 Unplanned interventions	6
2. Service scope	7
2.1 eHealth service	7
2.1.1 General	7
2.1.2 Abbreviations	8
2.2 Business criticality	8
2.3 Interdependencies	8
3. List of service levels	9
3.1 Detailed service level per service	10
3.1.1 Availability Consent Web App	10
3.1.2 Performance Consent Web App	11
3.1.3 Availability Consent Web Service	12
3.1.4 Performance Consent Web Service	13

To the attention of: "IT expert" willing to integrate this web service.



Service Level Agreement

Basic Service: Consent

Between

Service provider

eHealth Platform
Quai de Willebroeck, 38
1000 BRUSSELS

Service customer

User Community

To the attention of: the user community

Author: Service Management

Date: 04/04/2019

Version: 1.1

Status: Final

Type: Public

Confidentiality:

Language: English

Exhibit of: MSA



1. Document management

1.1 Document history

Version	Date	Author	Description of changes / remarks
2016.01	July, 2016	eHealth Service Management	First version
2018.01	February, 2018	eHealth Service Management	Modification KPI
1.1	06/08/2018	eHealth Service Management	Corrections
1.2	04/04/2019	eHealth Service Management	Modification KPI

1.2 Document references

Version	Title	Version	Date	Author
2	Master Service Agreement	2.0	22/11/2012	

1.3 Purpose of the document

The objective of this document is to define the Service Level Agreement for the set of services included in the [Basic Service Consent](#) proposed by the eHealth-platform. It defines the minimum level of service offered on the eHealth-platform, and provides eHealth's own understanding of service level offering, its measurement methods and its objectives in the end.

1.4 Features

The Consent Basic Service of the eHealth platform allows the handling of the patient's agreement to exchange his own medical data through authorised professionals and health care institutions.

This service is linked to the Metahub Service (see corresponding SLA for more information).

Consent Base Service is composed of two parts:

a) Consent Web Service (WS)

When a **"patient consent"** is active at the eHealth national level then a patient agrees to take part in the digital information exchange of his own medical data. This WS connects to the same database as the Metahub WS to manage and consult consents.

Only authorised types of professionals (doctor, nurse, dentist, midwife, and pharmacist) or professional organisations (Insurance organisations, CIN, Vitalink) may access the Consent WS.

Patients can also access it through specialized frontend software. They need to have a valid token from the eHealth STS to get access to this WS.

The two services covered by the WS are:

- *Get the patient consent [GetPatientConsentRequest]:* allows a hub to check the existence of an informed patient consent.
- *Declare or revoke patient consent [PutPatientConsentRequest - RevokePatientConsentRequest]:* declares/revokes an informed consent of a patient.



b) Consent Web Application

The Consent WA is not an exact representation of the functionality present in the Consent WS.

The three covered functionalities are:

- The informed patient consent: When a patient consent is active at the MetaHub level, the medical data of the patient can be shared between healthcare actors.
- The Therapeutic links: If a patient consent is active at the MetaHub level, healthcare professionals can access the medical documents of a patient only when a therapeutic link that justifies this consultation exists.
- If a Therapeutic exclusion exists between a patient and a healthcare professional, then this healthcare professional and institutions will not have access to the documents even if they fulfill all the requirements.

Note: the validity of the SSIN and support card numbers is checked through the ID Support WS, which relays the request to a CBSS WS at the consultation, the declaration and the revocation of the Therapeutic relations.

A patient can manage (get, put and revoke) through this WA:

- His/her consent and the corresponding audit
- His/her exclusions with one or more professionals and the corresponding audit
- His/her therapeutic links and the corresponding audit

A professional can manage for a certain patient:

- The patient's consent and the corresponding audit
- The patient's therapeutic links and the corresponding audit (only when certain pre-conditions are fulfilled)

1.5 Validity of the agreement

This document is valid as long as the *Basic Service Consent* is part of the eHealth-platform offering services.

Once a year, the levels of service proposed, will be reviewed and confirmed for the next year.

1.6 Service and maintenance window

1.6.1 Service window

The period during which the eHealth services are offered to the client applications, is defined in terms of days and hours. Standard working days are all days of the year, except during the biannual maintenance periods.

The following table summarises the eHealth service window.

		Service Window						
		Day of the week (closing days of Service Provider = Sunday)						
		Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday
Day period	00:00 – 07:00							
	07:00 – 08:00							
	08:00 – 16:30							
	16:30 – 19:00							
	19:00 – 20:00							
	20:00 – 24:00							



Legend	
	Timeslots where the Service must be available according to the SLA and where corrective actions will be taken to resolve detected Incidents.
	Timeslots where the Service will be available provided there are no blocking Incidents. If these incidents do appear, no corrective action will be taken.
	Timeslots where unavailability can occur.

1.6.2 Support window

Support Window								
		Day of the week (Closing days of Service Provider = Sunday)						
		Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday
Day period	00:00 – 07:00							
	07:00 – 08:00							
	08:00 – 16:30							
	16:30 – 19:00							
	19:00 – 20:00							
	20:00 – 24:00							

Legend	
	Timeslots for which the eHealth Call Center is available for the End-Users with a second line support for Infrastructure (HW, OS, Middleware and DB)
	Timeslots for which the eHealth Call Center is available for the End-Users with a second line support, including Application Support
	Timeslots for which the eHealth Call Center is unavailable for the End-Users. The End-User will have the possibility to record a voice message that will be treated on the next Workday.

1.6.3 Maintenance windows & planned interventions

The eHealth platform will strive for limiting as much as possible the impact and duration of the planned interventions. Today, the eHealth platform is committed to make efforts so planned unavailability's do not exceed one to a few hours per year. In case of maintenance requiring support from users, or affecting them, eHealth will notify them at least one week ahead.

1.6.4 Unplanned interventions


Under exceptional circumstances, unplanned interventions may be necessary in order to restore the service.



2. Service scope

2.1 eHealth service

2.1.1 General



The main components included in this SLA are:

- eHealth Consent WA (used by the patient and HC professionals on behalf of the patient, after access rights checks):
 - Informed Patient Consent management (Get/Declare/Revoke), including history data (using Get Patient Audit Trail method)
 - Therapeutic Links management (Get/Declare(Put)/Revoke) , including history data (some legal restrictions exist for HC Professionals) (using Get Therapeutic Links method)
 - Therapeutic Exclusion management (Get/Declare/Revoke) , including history data (only possible by the patient self) (using Get Patient Audit Trail method)

The declaration of a therapeutic exclusion or therapeutic link can be done by using the NIHI number of the HC professional or by searching him from a list built after filling (partially) name/first name of the HC professional.

The revocation of a therapeutic exclusion or link is done by selecting it in the list of the existing ones.

- eHealth Consent WS (used by the patient and by HC professionals and other authorized organisations on behalf of the patient, after access rights checks):
 - Informed Patient Consent management (Get/Declare/Revoke), including history data (using Get Patient Audit Trail method)



2.1.2 Abbreviations

AA	Attribute Authority
CBSS	Crossroads Bank for Social Security
CIN (NIC)	Collège Intermutualiste National
CoBRHA	Common Base Registry for Healthcare Actors
GMD	Global Medical Dossier
HC	Health Care
NIHII	National Institute for Health and Invalidation Insurance (INAMI / RIZIV)
SAMM	Source Authentique Migrée Mirrorée
SSIN	Social Security Identification Number
STS	Secure Token Service
UAM	User and Access Management

2.2 Business criticality

The business criticality of Consent service is **Gold** as it supports mandatory business processes processed synchronously and within some legal periods.

2.3 Interdependencies

N/A

3. List of service levels

Service	KPI	SL ID	Condition	Measure based on	Limit	Service Window	Objective Committed	Objective Target
Consent	Availability Consent WA		Test script passes	Fictitious request		Mo – Su 0:00 – 24:00	99,5%	99,9%
	Performance Consent WA (All)		Response time < 4 sec	Real transactions		Mo – Su 0:00 – 24:00	N/A	98,0%
	Availability Consent WS		Test script passes	Fictitious request		Mo – Su 0:00 – 24:00	99,5%	99,9%
	Performance Consent WS (All)		Response time < 4 sec	Real transactions		Mo – Su 0:00 – 24:00	N/A	99,0%
	Performance Consent WS - PutPatientConsent		Response time < 4 sec	Real transactions		Mo – Su 0:00 – 24:00	N/A	99,0%

Table 1: List of key performance indicators (KPI) per service



3.1 Detailed service level per service

3.1.1 Availability Consent Web App

Objectives				
Definition	<ul style="list-style-type: none"> The eHealth WA Consent is considered to be available when a test user can get: <ul style="list-style-type: none"> his Consent his therapeutic links his therapeutic link exclusions Planned interventions executed within the maintenance window are not recorded as unavailable time. 			
Measuring method	<ul style="list-style-type: none"> The availability of the different functionalities is measured by executing the test scripts every 5 minutes. When the script is executed with as result a status "OK", the test "passed". When the script is executed with another result, the test "failed". Measuring is always done on test scenarios 			
Calculation	$Availability = \frac{\sum Passed\ Tests \times 100}{\sum Total\ Tests} \%$ <ul style="list-style-type: none"> Total Tests = Total number of tests launched within corrected timeframe Passed Tests = Total number of tests that resulted in a status "OK" within the same timeframe Corrections are applicable on tests that are not taken into account because they were caused: <ul style="list-style-type: none"> by a Validated Authentic Source or partner application out of scope of this SLA by a failing monitoring tool 			
Reporting and evaluation period	<ul style="list-style-type: none"> The availability is calculated and reported monthly. Corrective interventions are initiated when appropriate. The formal evaluation however is done on a yearly basis. 			
Service Level Objectives	Functionality	Service Window	Service Level Objective	
			Committed	Target
	Availability Consent WA	Mo – Su 0:00 – 24:00	99,5%	99,9%



3.1.2 Performance Consent WA

Objectives				
Definition	<ul style="list-style-type: none"> The performance of the eHealth Consent WA refers to its response time. Response time meaning the time needed to execute a request. This request can be <ul style="list-style-type: none"> Get Patient Consent Declare/Revoke Patient Consent Get Therapeutic exclusions Declare/Revoke Therapeutic exclusions (depends on SAMM)/CoBRHA) Get Therapeutic links (depends on CIN) Declare/Revoke Therapeutic links (depends on SAMM/CoBRHA and CIN) Attention: The response time does not include: <ul style="list-style-type: none"> The time needed to deliver the information over the Internet The time needed to process the information at the end users premises. 			
Measuring method	<ul style="list-style-type: none"> This response time is measured on the Reverse Proxies. Both start time (request received) and stop time (answer sent to the End User) are measured and stored in a database. Measuring is done on real transactions, and only on those having a “stop time” within the measuring period. 			
Calculation	<ul style="list-style-type: none"> All response times are calculated: Stop time – Start time for every request. The percentage that meets the target is calculated based on following formula: $Performance = \frac{\sum Tests\ meeting\ the\ target \times 100}{\sum Total\ Tests} \%$			
Reporting and evaluation period	<ul style="list-style-type: none"> The performance is calculated and reported monthly. Corrective interventions are initiated when appropriate. The formal evaluation however is done on a yearly basis. 			
Service Level Objectives	Functionality	Target	Service Level Objective	
			Committed	Target
	Performance Consent WA	< 4 sec	N/A*	98%

*Due to external dependencies



3.1.3 Availability Consent Web Service

Objectives				
Definition	<ul style="list-style-type: none"> The eHealth WS Consent is considered to be available when it is reachable via the BUS and when the DBs are up and running (get request and evaluation of the response - keep Alive Test) Planned interventions executed within the Maintenance Window are not recorded as unavailable time. 			
Measuring method	<ul style="list-style-type: none"> The availability of the different functionalities is measured by executing the test scripts every 5 minutes. When the script is executed with as result a Status "OK", the test "passed". When the script is executed with an other result, the test "failed" Measuring is always done on test scenarios 			
Calculation	$Availability = \frac{\sum Passed\ Tests \times 100}{\sum Total\ Tests} \%$ <ul style="list-style-type: none"> Total Tests = Total number of tests launched within corrected timeframe Passed Tests = Total number of tests that resulted in a status "OK" within the same timeframe Corrections are applicable on tests that are not taken into account because they were caused: <ul style="list-style-type: none"> by a Validated Authentic Source or partner application out of scope of this SLA by a failing monitoring tool 			
Reporting and evaluation period	<ul style="list-style-type: none"> The availability is calculated and reported monthly. Corrective interventions are initiated when appropriate. The formal evaluation however is done on a yearly basis. 			
Service Level Objectives	Functionality	Service Window	Service Level Objective	
			Committed	Target
	Availability Consent WS	Mo – Su 0:00 – 24:00	99,5%	99,9%



3.1.4 Performance Consent Web Service

Objectives				
Definition	<ul style="list-style-type: none"> The performance of the eHealth Consent webservice refers to its response time. Response time meaning the time needed to execute a request. This request can be <ul style="list-style-type: none"> Get Patient Consent Put Patient Consent Revoke Patient Consent Attention: The response time does not include: <ul style="list-style-type: none"> The time needed to deliver the information over the Internet The time needed to process the information at the End Users premises. 			
Measuring method	<ul style="list-style-type: none"> This response time is measured on the Reverse Proxies. Both start time (request received) and stop time (answer sent to the End User) are measured and stored in a database. Measuring is done on real transactions, and only on those having a “stop time” within the measuring period. 			
Calculation	<ul style="list-style-type: none"> All response times are calculated: Stop time – Start time for every request. The percentage that meets the target is calculated based on following formula: $Performance = \frac{\sum Tests\ meeting\ the\ target \times 100}{\sum Total\ Tests} \%$			
Reporting and evaluation period	<ul style="list-style-type: none"> The performance is calculated and reported monthly. Corrective interventions are initiated when appropriate. The formal evaluation however is done on a yearly basis. 			
Service Level Objectives	Functionality	Target	Service Level Objective	
			Committed	Target
	Performance Consent WS (all)	< 4 sec	N/A*	99,0%
	Performance Consent WS PutPatientConsent	< 4 sec	N/A*	99,0%

*Due to external dependencies

